

IV Glove 2.0 Adult

Pre-application: Clean the site per hospital protocol, and ensure the site is dry before applying the

- 1. Follow standard IV insertion as per hospital protocol 2. Cover the insertion site with
- medical tape or Tegaderm
- 3. Open the transparent covered dome of the IV Glove by pulling up on the cover
- 4. Place 3 to 4 of patient's fingers through the IV Glove finger loop according to patient and clinician preference
- 5. Fasten the wrist clip and adjust the blue stretchy wrist bands to ensure comfort and securement.
- 6. Ensure that you can slide your index and middle finger under the strap to ensure the wrist strap is not too tiaht
- 7. If necessary, adjust the finger loop straps to tighten by pulling the ends of the finger loop straps through the top of the device
- 8. Ensure capillary refill of less than 2 seconds distal to insertion site
- 9. Place catheter tubing into patented S-Clip design
- 10. Carefully close the protective dome, ensuring the pins 'click' in place
- 11. IV Glove is ready for use



for use

IV Glove 2.0 Adult

Assessment

- -Follow hospital protocol for assessment
- -Simply inspect through the transparent covered dome to ensure proper catheter placement (no redness, swelling, etc.)
- -If further inspection is required, simply lift the transparent covered dome for closer inspection

Removal

- 1. Open the transparent covered dome and remove the tubing from the S-Clip 2. Loosen the wrist strap by removing
- 2.Loosen the wrist strap by removing the pin from the hole
- 3.Loosen the finger loop by removing the pin from the hole
- 4.Slide the IV Glove off the patient's hand, taking care to mind the IV insertion site
- 5.Remove the medical tape or Tegaderm
- 6.Remove the IV catheter

Post-application

- -IV Glove is made of 100% recyclable materials
- -Dispose of the IV Glove per hospital protocol

Scan for instruction video





